

Ecto-5'-nucleotidase, adenosine and transmembrane adenylyl cyclase signalling regulate basal carotid body chemoafferent outflow and establish the sensitivity to hypercapnia

Holmes, Andrew P; Nunes, Ana Rita; Cann, Martin J; Kumar, Prem

DOI:

[10.1007/978-3-319-18440-1_32](https://doi.org/10.1007/978-3-319-18440-1_32)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Holmes, AP, Nunes, AR, Cann, MJ & Kumar, P 2015, Ecto-5'-nucleotidase, adenosine and transmembrane adenylyl cyclase signalling regulate basal carotid body chemoafferent outflow and establish the sensitivity to hypercapnia. in C Peers, P Kumar, C Wyatt, E Gauda, CA Nurse & N Prabhakar (eds), *Arterial Chemoreceptors in Physiology and Pathophysiology*. vol. 860, Advances in Experimental Medicine and Biology, vol. 860, Springer, pp. 279-289. https://doi.org/10.1007/978-3-319-18440-1_32

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Download date: 05. May. 2023

Ecto-5'-nucleotidase, adenosine and transmembrane adenylyl cyclase signalling regulate basal carotid body chemoafferent outflow and establish the sensitivity to hypercapnia.

Andrew P. Holmes¹, Ana Rita Nunes², Martin J. Cann³ & Prem Kumar¹

¹. School of Clinical and Experimental Medicine, University of Birmingham, Edgbaston, Birmingham, UK, B15 2TT

². CEDOC, Chronic Diseases Research Center, NOVA Medical School / Faculdade de Ciências Médicas, Universidade Nova de Lisboa, Campo Mártires da Pátria, 130, 1169-056 Lisboa, Portugal

³. School of Biological and Biomedical Sciences, South Road, Durham University, Durham, UK, DH1 3LE

Key Words: carotid body, adenosine, hypercapnia, ecto-5'-nucleotidase, cAMP

Word count: 2670

Corresponding Author: Andrew P. Holmes

Address: Level 1 IBR, School of Clinical and Experimental Medicine, University of Birmingham, Edgbaston, Birmingham, UK, B15 2TT

Email: a.p.holmes@bham.ac.uk

Tel: 0121 414 46938

Abstract

Carotid body (CB) stimulation by hypercapnia causes a reflex increase in ventilation and, along with the central chemoreceptors, this prevents a potentially lethal systemic acidosis. Control over the CB chemoafferent output during normocapnia and hypercapnia most likely involves multiple neurotransmitters and neuromodulators including ATP, acetylcholine, dopamine, serotonin and adenosine, but the precise role of each is yet to be fully established. In the present study, recordings of chemoafferent discharge frequency were made from the isolated *in vitro* CB in order to determine the contribution of adenosine, derived specifically from extracellular catabolism of ATP, in mediating basal chemoafferent activity and responses to hypercapnia. Pharmacological inhibition of ecto-5'-nucleotidase (CD73), a key enzyme required for extracellular generation of adenosine from ATP, using α,β -methylene ADP, virtually abolished the basal normocapnic single fibre discharge frequency (superfusate $PO_2 \sim 300$ mmHg, $PCO_2 \sim 40$ mmHg) and diminished the chemoafferent response to hypercapnia ($PCO_2 \sim 80$ mmHg). These effects were mimicked by the blockade of adenosine receptors with 8-(p-sulfophenyl) theophylline. The excitatory impact of adenosinergic signalling on CB hypercapnic sensitivity is most likely to be conferred through changes in cAMP. Here, inhibition of transmembrane, but not soluble, adenylate cyclases, by SQ22536 and KH7 respectively, produced a rapid reduction in normocapnic single fibre activity and inhibited the elevation evoked by hypercapnia by approximately 50%. These data therefore identify a functional role for CD73 derived adenosine and transmembrane adenylate cyclases, in modulating the basal chemoafferent discharge frequency and in priming the CB to hypercapnic stimulation.

1 Introduction

The mammalian carotid bodies (CBs) are the primary peripheral chemoreceptors that respond to acute hypoxia, and stimulation drives the reflex hyperventilation that acts to preserve O₂ delivery and metabolism in the brain and vital organs (Kumar, 2009). However, the importance of the CB in the regulation of breathing is not limited to hypoxia. Continuous chemoafferent input from CB into the central nervous system (CNS) is thought to account for up to 60% of eupneic ventilation (Blain *et al.*, 2009). Furthermore, the CBs are activated by other blood stimuli including hypercapnia and acidosis, both of which are potentially lethal if not adequately countered.

It was originally proposed that approximately 30-50% of the reflex ventilatory response to arterial hypercapnia was mediated through direct stimulation of the CB (Heeringa *et al.*, 1979; Rodman *et al.*, 2001), with the remaining contribution arising from chemoreceptors located in the CNS (Nattie, 1999). However, a more recent investigation has shown that a complete silencing of the CB chemoafferent output significantly depresses the central CO₂ chemoreceptor sensitivity (Blain *et al.*, 2010). Therefore, in addition to direct hypercapnic CB excitation, maintenance of a tonic CB chemoafferent signal into the CNS seems to have an important role in establishing the hypercapnic sensitivity of the central chemoreceptors and further emphasises the importance of the CB in blood CO₂ and pH homeostasis.

Control over the CB chemoafferent output in basal conditions and during hypercapnia most likely involves multiple neurotransmitters and neuromodulators including ATP, acetylcholine, dopamine, serotonin and adenosine (Nurse, 2010), but the contribution of each is yet to be fully characterised. Despite adenosine being an established neuromodulatory substance in the central nervous system (Cunha, 2001), its source and physiological role in the CB remains somewhat unresolved. Up to now the majority of investigations have focused on the action of adenosine in mediating the CB response to acute changes in O₂ tension (McQueen & Ribeiro, 1986; Conde *et al.*, 2006; Conde *et al.*, 2012) or the adaptations following chronic hypoxia (Livermore & Nurse, 2013). However, the physiological importance and source of endogenous adenosine in establishing basal CB chemoafferent activity and responses to hypercapnia is unclear.

Physiologically active concentrations of adenosine, present in the synapse between the CB type I cell and the adjacent chemoafferent neurone, may originate from multiple sources. First, adenosine may be generated via extracellular catabolism of ATP; an excitatory neurotransmitter that is secreted from type I and type II cells (Zhang *et al.*, 2012; Piskuric & Nurse, 2013). This conversion requires two key enzymes; ectonucleoside triphosphate diphosphohydrolyase 1 (CD39) and ecto-5'-nucleotidase (CD73) (Bianchi & Spychala, 2003). Second, adenosine may be formed in the type I cell and then released into the synapse through the bidirectional equilibrative nucleotide transporter (ENT) (Cass *et al.*, 1998). Measurements of whole organ adenosine suggest a possible contribution of both sources to the total 'pool' of available adenosine and this may be subject to variation depending on the ambient level of O₂ (Conde & Monteiro, 2004; Conde *et al.*, 2012). This current study aims to more clearly define the

functional importance of endogenous adenosine specifically generated from extracellular catabolism of ATP in modulating the sensory neuronal output of the CB, both under basal conditions and during hypercapnia.

The four G-protein coupled adenosine receptors cloned to date (A_1 , A_{2A} , A_{2B} and A_3) all exert their actions through inhibition or excitation of transmembrane adenylyl cyclases (tmACs) and production of cAMP (reviewed in (Ribeiro & Sebastiao, 2010)). Thus any functional role of adenosine is likely to be conferred through modifications in cAMP accumulation. We have previously reported that the CB expresses numerous different tmAC mRNA transcripts (Nunes *et al.*, 2013) as well as the soluble AC (sAC) isoform (Nunes *et al.*, 2009). CB cAMP is elevated during hypercapnia (Perez-Garcia *et al.*, 1990) and, in isolated type I cells, cAMP analogues potentiate inward Ca^{2+} current in a manner that is quantitatively similar to hypercapnia (Summers *et al.*, 2002). Here we investigate whether direct targeting of tmAC or sAC impairs the CB chemoafferent outflow in normocapnia and/or hypercapnia.

2 Methods

2.1 Ethical approval

All surgical procedures were performed in accordance with project and personal licences issued under the UK Animals (Scientific Procedures) Act 1986 and were approved by the Biomedical Services Unit at the University of Birmingham.

2.2 Extracellular recordings of single and few-fibre chemoafferent neurones

The whole carotid bifurcation along with the attached carotid sinus nerve (CSN) and CB were isolated from adult male Wistar rats (50–200 g) under inhalation anaesthesia (2–4% isoflurane in O_2). Following tissue removal, animals were immediately killed by exsanguination. The bifurcations were rapidly transferred into an ice cold bicarbonate buffered extracellular Krebs solution containing, in mM: 115 NaCl, 4.5 KCl, 1.25 NaH_2PO_4 , 5 Na_2SO_4 , 1.3 $MgSO_4$, 24 $NaHCO_3$, 2.4 $CaCl_2$, 11 D-glucose, equilibrated with 95% O_2 and 5% CO_2 .

Extracellular recordings of single or few-fibre chemoafferent activity were made from the cut end of the CSN using glass suction electrodes pulled from GC150-10 capillary glass (Harvard Apparatus, Edenbridge, UK). The chemoafferent derived voltage was recorded using a CED micro1401 and visualised on a PC with Spike2 software (Cambridge Electronic Design, Cambridge, UK). The chemoafferent voltage signal was sampled at 15000 Hz. Using the in-built wavemark analysis in the Spike2 software, electrical activity originating from a single chemoafferent fibre was determined by its unique 'wavemark' signature based on frequency, shape and amplitude.

2.3 Experimental solutions and drugs.

During experimentation, whole CBs were continuously superfused with a standard bicarbonate buffered Krebs solution containing, in mM: 115 NaCl, 4.5 KCl, 1.25 NaH_2PO_4 , 5 Na_2SO_4 , 1.3 $MgSO_4$, 24 $NaHCO_3$,

2.4 CaCl₂ and 11 D-glucose. All solutions were heated to 37°C using a water bath (Grant W14; Grant Instruments, Cambridge, UK) and had a pH of 7.4 except during hypercapnic acidosis.

Chemoafferent responses to hypercapnia acidosis were induced by raising the superfusate PCO₂ from approximately 40 mmHg (pH 7.4) to 80 mmHg (pH 7.15) at a constant PO₂ (300 mmHg), as has been previously reported (Bin-Jalilah *et al.*, 2005). Since the chemoafferent response to hypercapnia is thought to peak initially and then adapt to a lower sustained frequency (Black *et al.*, 1971), measurements of chemoafferent activity were taken from the fifth minute of the hypercapnic stimulus, after a relatively steady state frequency had been achieved. CO₂ sensitivity was subsequently calculated as the increase in single fibre discharge frequency per mmHg increase in superfusate PCO₂ (Δ Hz / mmHg PCO₂), given that the rise in discharge frequency is linear over this PCO₂ range (Biscoe *et al.*, 1970; Fitzgerald & Parks, 1971; Pepper *et al.*, 1995; Vidruk *et al.*, 2001).

2.4 Analysis of data

Values are expressed as mean \pm standard error of mean unless otherwise stated. Statistical analysis was performed using i) a paired 2-tailed student's t-test or ii) repeated measures one way Analysis of Variance (ANOVA) with Bonferroni or Dunnett's post hoc analysis where appropriate (Prism v5; GraphPad Software, La Jolle, CA, USA). Significance was taken as $p < 0.05$.

3 Results

3.1 Basal chemoafferent outflow is dependent on ecto-5'-nucleotidase (CD73) activity and adenosinergic signalling.

Basal chemoafferent output from the CB provides the peripheral component of the drive to breathe. Initial experiments sought to determine whether or not adenosine generated from extracellular ATP catabolism has any part in establishing this sensory neuronal activity of the CB. Pharmacological targeting of ecto-5'-nucleotidase (CD73; a key membrane bound enzyme involved in the formation of adenosine from extracellular ATP) using the inhibitor α,β -methylene ADP (AOPCP; 100 μ M; Conde & Monteiro, 2004) had a striking effect on basal single fibre activity, diminishing the discharge frequency by $91.2 \pm 3.8\%$ (Fig. 1a *upper* & 1b). A similar ($93.9 \pm 1.9\%$) reduction in basal frequency was observed in the presence of the adenosine receptor antagonist 8-(p-sulfophenyl) theophylline (8-SPT, 300 μ M; Wyatt *et al.*, 2007) (Fig. 1a *middle* & 1b). The four G-protein coupled adenosine receptors cloned to date (A₁, A_{2A}, A_{2B} and A₃) all exert their actions through inhibition or excitation of transmembrane adenylyl cyclases (tmACs) and modification of cAMP production. In the current investigation, the presence of 9-(Tetrahydro-2-furanyl)-9H-purin-6-amine (SQ22536; a tmAC inhibitor; 200 μ M, IC₅₀ = 20 μ M; Rocher *et al.*, 2009) caused a $70.2 \pm 16.0\%$ decrease in basal single fibre frequency (Fig. 1a *lower* & 1b). This is consistent with the idea that tonic generation of adenosine from ATP modulates basal CB neuronal outflow through production of cAMP.

3.2 Chemoafferent responses to hypercapnia are dependent on ecto-5'-nucleotidase (CD73) activity, adenosinergic receptor stimulation and transmembrane adenylyl cyclase production of cAMP.

Experiments were designed to establish or rule out a potential role for endogenous adenosine in mediating the heightened sensory neuronal activity of the CB in hypercapnia. Selective inhibition of CD73 with AOPCP (100 μ M) diminished the single fibre discharge frequency recorded in both normocapnic (40 mmHg PCO₂) and hypercapnic (80 mmHg PCO₂) conditions (Fig. 2a & b). Furthermore, AOPCP caused a dramatic ($98.0 \pm 1.8\%$) reduction in CB CO₂ sensitivity (Fig. 2c). This inhibitory effect was rapidly reversed once the drug was removed from the superfusate as demonstrated in the raw trace example in Fig. 2a. Similar observations were made in the presence of 8-SPT (300 μ M). 8-SPT significantly attenuated chemoafferent activity in both normocapnia and hypercapnia (Fig. 2d) and markedly decreased the CB CO₂ sensitivity by $81.5 \pm 5.8\%$ (Fig. 2e).

As with the basal discharge, the excitatory impact of adenosinergic signalling on CB hypercapnic sensitivity is most likely to be conferred through activation of tmACs coupled to the adenosine receptors. Here, addition of the tmAC inhibitor SQ22536 (200 μ M), produced a rapid reduction in normocapnic single fibre activity and inhibited the elevation evoked by hypercapnia (Fig. 3a & b). In every fibre tested the absolute increase in hypercapnic discharge was reduced in the presence of SQ22536. Accordingly, further analysis showed that SQ22536 (200 μ M) produced a $47.1 \pm 9.4\%$ decline in CB CO₂ sensitivity (Fig. 3c). Normal hypercapnic responses were readily restored following removal of the agent from the superfusate, as exemplified in Fig. 3a. In contrast, the soluble adenylyl cyclase (sAC) antagonist, 2-(1H-Benzo[d]imidazol-2-ylthio)-N'-(5-bromo-2-hydroxybenzylidene)propanehydrazide (KH7; 10 μ M; Nunes *et al.*, 2013), failed to impact on basal chemoafferent activity or the response to hypercapnia (Fig. 3d & e). KH7 (10 μ M) did not alter CB CO₂ sensitivity even after prolonged incubation (Fig. 3d & f). These data therefore suggest that cAMP production mediates a component of the functional chemoafferent response to hypercapnia and that this is selectively dependent on tmAC rather than sAC activation.

4 Discussion

4.1 CD73, adenosine and tmAC signalling mediate basal CB activity and the sensitivity to hypercapnia.

The findings presented in this study indicate that extracellular adenosine formed selectively through catabolism of ATP, in the presence of ecto-5'-nucleotidase (CD73), has an important neuromodulatory role in mediating the CB sensory neuronal discharge, both under basal conditions and during stimulation by hypercapnia. In addition we show that tmAC but not sAC activity is necessary for full expression of CB hypercapnic sensitivity.

It has been previously reported that inhibition of ATP metabolism in normoxia by AOPCP decreases CB adenosine content (Conde & Monteiro, 2004). Our observations suggest that this source of adenosine is functionally required in order to generate the basal chemoafferent outflow. The impact of adenosine is

most likely to be conferred through modifications in cAMP, given that the A_{2A} and A_{2B} adenosine receptor subtypes expressed in the CB (Gauda *et al.*, 2000; Kobayashi *et al.*, 2000; Conde *et al.*, 2006) are coupled to tmACs (Ribeiro & Sebastiao, 2010). Accordingly, in this investigation pharmacological inhibition of tmACs by SQ22536 caused a significant reduction in basal chemoafferent activity.

The basal sensory output from the CB accounts for up to 60% of the drive to breath at rest (Blain *et al.*, 2009) and may contribute to the resting sympathetic outflow to the vasculature (Kumar & Prabhakar, 2012). Furthermore, it is the chronic rise in basal CB activity following CIH that is thought to drive the hypertension in patients with OSA (Narkiewicz *et al.*, 1998; Peng *et al.*, 2003). It is probable that overall chemoafferent output in normoxia is dependent on both spontaneous pre-synaptic (type I cell) depolarisation and neurotransmitter release, along with subsequent post-synaptic receptor activation. Chemoafferent neurones may also generate a degree of spontaneous activity. In large clusters of type I cells, spontaneous cellular depolarisation is reported to be a consequence of an attenuation of the background K⁺ current, which is itself regulated by local 5-HT and PKC activation (Zhang & Nurse, 2000; Zhang *et al.*, 2003). The observations presented here suggest an additional signalling mechanism that is dependent on adenosine formation and tmAC mediated cAMP production.

It is becoming more apparent that the CB has a principal role in countering rises in arterial CO₂. As well as direct stimulation of the CB accounting for approximately 30-50% of the reflex hypercapnic ventilatory response (Heeringa *et al.*, 1979; Rodman *et al.*, 2001), it has been recently shown that the CB chemoafferent outflow is necessary in establishing the CO₂ sensitivity of the central medullary chemoreceptors (Blain *et al.*, 2010). Despite these important findings, the full transduction mechanisms that lead to an increase in CB discharge frequency during hypercapnia are still elusive. The results described here provide evidence that the heightened chemoafferent activity in hypercapnia is dependent on CD73 mediated catabolism of ATP to form adenosine. The principal origin of the synaptic ATP is most likely to be the type I cell given that ATP is secreted as a neurotransmitter in both normocapnic (Conde *et al.*, 2012) and hypercapnic conditions (Zhang & Nurse, 2004). Interestingly, in chemosensitive regions of the brainstem, hypercapnia-evoked ATP release occurs through the gap junctional protein, connexin26 (Cx26) (Huckstepp *et al.*, 2010b). Direct binding of CO₂ prevents pore occlusion and effectively 'traps' Cx26 in an open conformation (Huckstepp *et al.*, 2010a; Meigh *et al.*, 2013). In the CB, recent evidence has identified a mechanism of ATP induced ATP release through pannexin channels in type II cells (Zhang *et al.*, 2012). Whether ATP is released from type II cells in normoxia and/or hypercapnia, thus acting as a substrate for additional adenosine formation, could be an interesting area for future investigating.

We also demonstrate that inhibitory targeting of tmACs also reduced the CO₂ sensitivity of the CB. This is in agreement with the idea that adenosine partially establishes the chemoafferent sensitivity of the CB to hypercapnia through tmAC dependent cAMP production. In the later instance however, a considerable proportion of the response to hypercapnia is preserved, which was in contrast to the results described using 8-SPT and AOPCP. This may have been due to incomplete run-down in tmAC function

using SQ22536 or, more likely, that adenosine has additional actions that are independent of cAMP. These alternative downstream signalling pathways possibly involve activation of PKC or phospholipase C, as has been proposed for a component of the excitatory adenosine neuromodulation in the CNS (Cunha, 2001).

The finding that SQ22536 but not KH7 reduces chemoafferent excitation in hypercapnia, suggests that cAMP regulation of CB CO₂ sensitivity is itself determined by tmAC and not sAC enzymatic activity. Although sAC is present in the CB (Nunes *et al.*, 2009) and is directly stimulated by binding of CO₂ or HCO₃⁻ (Townsend *et al.*, 2009), it appears to have no functional contribution in moderating the chemoafferent frequency in hypercapnia. This is in accordance with our previous findings where KH7 failed to modify CB cAMP content when the organ was exposed to isohydric hypercapnia (Nunes *et al.*, 2013). Thus we propose that rises in CO₂ or intracellular HCO₃⁻ alone are insufficient to stimulate sAC enough to have functional effects on chemoafferent discharge in hypercapnia.

4.3 Conclusions

Endogenous adenosine produced from extracellular catabolism of ATP in the presence of ecto-5'-nucleotidase (CD73) is necessary for the generation of a basal chemoafferent discharge frequency. This basal discharge may be of clinical significance in a number of cardiorespiratory disorders where enhanced CB activity is associated with increased sympathetic outflow and thus with potential increase in patient morbidity and mortality. In addition this source of adenosine and tmAC generation of cAMP, acting downstream of adenosine receptors, are required for the full expression of the CB sensitivity to hypercapnia.

References

- Bianchi V & Spychala J. (2003). Mammalian 5'-nucleotidases. *J Biol Chem* **278**, 46195-46198.
- Bin-Jaliah I, Maskell PD & Kumar P. (2005). Carbon dioxide sensitivity during hypoglycaemia-induced, elevated metabolism in the anaesthetized rat. *J Physiol* **563**, 883-893.
- Biscoe TJ, Purves MJ & Sampson SR. (1970). The frequency of nerve impulses in single carotid body chemoreceptor afferent fibres recorded in vivo with intact circulation. *J Physiol* **208**, 121-131.
- Black AM, McCloskey DI & Torrance RW. (1971). The responses of carotid body chemoreceptors in the cat to sudden changes of hypercapnic and hypoxic stimuli. *Respir Physiol* **13**, 36-49.
- Blain GM, Smith CA, Henderson KS & Dempsey JA. (2009). Contribution of the carotid body chemoreceptors to eupneic ventilation in the intact, unanesthetized dog. *J Appl Physiol* **106**, 1564-1573.
- Blain GM, Smith CA, Henderson KS & Dempsey JA. (2010). Peripheral chemoreceptors determine the respiratory sensitivity of central chemoreceptors to CO₂. *J Physiol* **588**, 2455-2471.
- Cass CE, Young JD & Baldwin SA. (1998). Recent advances in the molecular biology of nucleoside transporters of mammalian cells. *Biochem Cell Biol* **76**, 761-770.
- Conde SV & Monteiro EC. (2004). Hypoxia induces adenosine release from the rat carotid body. *J Neurochem* **89**, 1148-1156.
- Conde SV, Monteiro EC, Rigual R, Obeso A & Gonzalez C. (2012). Hypoxic intensity: a determinant for the contribution of ATP and adenosine to the genesis of carotid body chemosensory activity. *J Appl Physiol* **112**, 2002-2010.
- Conde SV, Obeso A, Vicario I, Rigual R, Rocher A & Gonzalez C. (2006). Caffeine inhibition of rat carotid body chemoreceptors is mediated by A_{2A} and A_{2B} adenosine receptors. *J Neurochem* **98**, 616-628.
- Cunha RA. (2001). Adenosine as a neuromodulator and as a homeostatic regulator in the nervous system: different roles, different sources and different receptors. *Neurochem Int* **38**, 107-125.
- Fitzgerald RS & Parks DC. (1971). Effect of hypoxia on carotid chemoreceptor response to carbon dioxide in cats. *Respir Physiol* **12**, 218-229.
- Gauda EB, Northington FJ, Linden J & Rosin DL. (2000). Differential expression of A_{2A}, A₁-adenosine and D₂-dopamine receptor genes in rat peripheral arterial chemoreceptors during postnatal development. *Brain Res* **872**, 1-10.
- Heeringa J, Berkenbosch A, de Goede J & Olivier CN. (1979). Relative contribution of central and peripheral chemoreceptors to the ventilatory response to CO₂ during hyperoxia. *Respir Physiol* **37**, 365-379.
- Huckstepp RT, Eason R, Sachdev A & Dale N. (2010a). CO₂-dependent opening of connexin 26 and related beta connexins. *J Physiol* **588**, 3921-3931.
- Huckstepp RT, id Bihi R, Eason R, Spyer KM, Dicke N, Willecke K, Marina N, Gourine AV & Dale N. (2010b). Connexin hemichannel-mediated CO₂-dependent release of ATP in the medulla oblongata contributes to central respiratory chemosensitivity. *J Physiol* **588**, 3901-3920.
- Kobayashi S, Conforti L & Millhorn DE. (2000). Gene expression and function of adenosine A_{2A} receptor in the rat carotid body. *Am J Physiol Lung Cell Mol Physiol* **279**, L273-282.

- Kumar P. (2009). Systemic effects resulting from carotid body stimulation-invited article. *Adv Exp Med Biol* **648**, 223-233.
- Kumar P & Prabhakar NR. (2012). Peripheral Chemoreceptors: Function and Plasticity of the Carotid Body. *Comprehensive Physiology*, 141–219.
- Livermore S & Nurse CA. (2013). Enhanced adenosine A2b receptor signaling facilitates stimulus-induced catecholamine secretion in chronically hypoxic carotid body type I cells. *Am J Physiol Cell Physiol* **305**, C739-750.
- McQueen DS & Ribeiro JA. (1986). Pharmacological characterization of the receptor involved in chemoexcitation induced by adenosine. *Br J Pharmacol* **88**, 615-620.
- Meigh L, Greenhalgh SA, Rodgers TL, Cann MJ, Roper DI & Dale N. (2013). CO₂ directly modulates connexin 26 by formation of carbamate bridges between subunits. *Elife* **2**.
- Narkiewicz K, van de Borne PJH, Montano N, Dyken ME, Phillips BG & Somers VK. (1998). Contribution of tonic chemoreflex activation to sympathetic activity and blood pressure in patients with obstructive sleep apnea. *Circulation* **97**, 943-945.
- Nattie E. (1999). CO₂, brainstem chemoreceptors and breathing. *Prog Neurobiol* **59**, 299-331.
- Nunes AR, Holmes AP, Sample V, Kumar P, Cann MJ, Monteiro EC, Zhang J & Gauda EB. (2013). Bicarbonate-sensitive soluble and transmembrane adenylyl cyclases in peripheral chemoreceptors. *Respir Physiol Neurobiol* **188**, 83-93.
- Nunes AR, Monteiro EC, Johnson SM & Gauda EB. (2009). Bicarbonate-regulated soluble adenylyl cyclase (sAC) mRNA expression and activity in peripheral chemoreceptors. *Adv Exp Med Biol* **648**, 235-241.
- Nurse CA. (2010). Neurotransmitter and neuromodulatory mechanisms at peripheral arterial chemoreceptors. *Exp Physiol* **95**, 657-667.
- Peng YJ, Overholt JL, Kline D, Kumar GK & Prabhakar NR. (2003). Induction of sensory long-term facilitation in the carotid body by intermittent hypoxia: implications for recurrent apneas. *Proc Natl Acad Sci U S A* **100**, 10073-10078.
- Pepper DR, Landauer RC & Kumar P. (1995). Postnatal development of CO₂-O₂ interaction in the rat carotid body in vitro. *J Physiol* **485** (Pt 2), 531-541.
- Perez-Garcia MT, Almaraz L & Gonzalez C. (1990). Effects of different types of stimulation on cyclic AMP content in the rabbit carotid body: functional significance. *J Neurochem* **55**, 1287-1293.
- Piskuric NA & Nurse CA. (2013). Expanding role of ATP as a versatile messenger at carotid and aortic body chemoreceptors. *J Physiol* **591**, 415-422.
- Ribeiro JA & Sebastiao AM. (2010). Modulation and metamodulation of synapses by adenosine. *Acta Physiol (Oxf)* **199**, 161-169.
- Rocher A, Caceres AI, Almaraz L & Gonzalez C. (2009). EPAC signalling pathways are involved in low PO₂ chemoreception in carotid body chemoreceptor cells. *J Physiol* **587**, 4015-4027.
- Rodman JR, Curran AK, Henderson KS, Dempsey JA & Smith CA. (2001). Carotid body denervation in dogs: eupnea and the ventilatory response to hyperoxic hypercapnia. *J Appl Physiol* **91**, 328-335.
- Summers BA, Overholt JL & Prabhakar NR. (2002). CO(2) and pH independently modulate L-type Ca(2+) current in rabbit carotid body glomus cells. *J Neurophysiol* **88**, 604-612.

- Townsend PD, Holliday PM, Fenyk S, Hess KC, Gray MA, Hodgson DRW & Cann MJ. (2009). Stimulation of Mammalian G-protein-responsive Adenylyl Cyclases by Carbon Dioxide. *J Biol Chem* **284**, 784-791.
- Vidruk EH, Olson EB, Jr., Ling L & Mitchell GS. (2001). Responses of single-unit carotid body chemoreceptors in adult rats. *J Physiol* **531**, 165-170.
- Wyatt CN, Mustard KJ, Pearson SA, Dallas ML, Atkinson L, Kumar P, Peers C, Hardie DG & Evans AM. (2007). AMP-activated protein kinase mediates carotid body excitation by hypoxia. *J Biol Chem* **282**, 8092-8098.
- Zhang M, Fearon IM, Zhong H & Nurse CA. (2003). Presynaptic modulation of rat arterial chemoreceptor function by 5-HT: role of K⁺ channel inhibition via protein kinase C. *J Physiol* **551**, 825-842.
- Zhang M & Nurse CA. (2000). Does endogenous 5-HT mediate spontaneous rhythmic activity in chemoreceptor clusters of rat carotid body? *Brain Res* **872**, 199-203.
- Zhang M & Nurse CA. (2004). CO₂/pH chemosensory signaling in co-cultures of rat carotid body receptors and petrosal neurons: role of ATP and ACh. *J Neurophysiol* **92**, 3433-3445.
- Zhang M, Piskuric NA, Vollmer C & Nurse CA. (2012). P2Y₂ receptor activation opens pannexin-1 channels in rat carotid body type II cells: potential role in amplifying the neurotransmitter ATP. *J Physiol* **590**, 4335-4350.

Figure legends

Fig. 1 The basal chemoafferent discharge frequency of the carotid body (CB) is critically dependent on ecto-5'-nucleotidase (CD73) mediated formation of adenosine. **(a)** Example raw neuronal traces demonstrating the inhibitory impact of three different compounds on basal chemoafferent outflow: AOPCP (100 μ M; inhibitor of CD73), 8-SPT (300 μ M; non-selective adenosine receptor antagonist) and SQ22536 (200 μ M; inhibitor of transmembrane adenylate cyclases). For each trace, overdrawn action potentials are shown inset to exhibit single fibre discrimination. **(b)** Mean single fibre basal frequencies in the presence and absence of each pharmacological agent. Data presented is from 6 fibres (n=6) from 4 CB preparations (AOPCP), 6 fibres (n=6) from 4 CB preparations (8-SPT) and 5 fibres (n=5) from 4 CB preparations (SQ22536). Error bars indicate \pm S.E.M. * denotes $P < 0.05$ compared with control basal discharge frequency; paired Student's t-test.

Fig. 2 Carotid body (CB) responses to hypercapnia are mediated by adenosine generated from ecto-5'-nucleotidase (CD73). **(a)** An example electrophysiological recording of the CB response to 5 minutes of hypercapnia ($PCO_2 = 80$ mmHg) in the presence and absence of the CD73 inhibitor AOPCP. Raw discharge is shown (upper) along with frequency histograms (lower). Overdrawn action potentials are shown inset to demonstrate the single fibre discrimination used to measure frequency. The inhibitory action of AOPCP was readily reversible. **(b)** Mean discharge frequencies recorded in normocapnia ($PCO_2 = 40$ mmHg) and hypercapnia ($PCO_2 = 80$ mmHg), in control conditions and following addition of AOPCP. Error bars indicate \pm S.E.M. * denotes $P < 0.05$ compared with control group; one way repeated measures ANOVA with Bonferroni post hoc analysis. **(c)** Calculated mean CO_2 sensitivity (Δ Hz / mmHg PCO_2) in control conditions and following AOPCP drug application. Error bars indicate \pm S.E.M. * denotes $P < 0.05$ compared with control group; paired Student's t-test. For B) and C) data is from 6 fibres (n=6) from 5 CB preparations. **(d)** Mean discharge frequencies recorded in normocapnia and hypercapnia, in control conditions and in the presence of 8-SPT (300 μ M), the non-selective adenosine receptor antagonist. Error bars indicate \pm S.E.M. * denotes $P < 0.05$ compared with control group; one way repeated measures ANOVA with Bonferroni post hoc analysis. **(e)** Calculated mean CO_2 sensitivity (Δ Hz / mmHg PCO_2) for control and 8-SPT groups. Error bars indicate \pm S.E.M. * denotes $P < 0.05$ compared with control group; paired Student's t-test. For D) and E) data is from 7 fibres (n=7) from 3 CB preparations.

Fig. 3 Rises in chemoafferent activity in hypercapnia are dependent on transmembrane (tmAC) but not soluble adenylate cyclase (sAC) activity. **(a)** Characteristic example recording of the response to hypercapnia ($PCO_2 = 80$ mmHg) in the presence and absence of the tmAC inhibitor SQ22536 (200 μ M). Raw discharge is shown (upper) along with frequency histograms (lower). Overdrawn action potentials are shown inset to demonstrate the single fibre discrimination. The inhibitory impact of SQ22536 was fully reversible. **(b)** Mean single fibre discharge frequencies recorded in normocapnia ($PCO_2 = 40$

mmHg) and hypercapnia ($\text{PCO}_2 = 80 \text{ mmHg}$), in control conditions and following addition of SQ22536 to the superfusate. Error bars indicate \pm S.E.M. * denotes $P < 0.05$ compared with control group; one way repeated measures ANOVA with Bonferroni post hoc analysis. **(c)** Calculated mean CO_2 sensitivity ($\Delta \text{ Hz} / \text{ mmHg } \text{PCO}_2$) in control conditions during SQ22536 drug application. Error bars indicate \pm S.E.M. * denotes $P < 0.05$ compared with control group; paired Student's t-test. For (b) and (c) the data presented is from 7 fibres ($n=7$) from 4 CB preparations. **(d)** As for (a) but in the presence of the sAC inhibitor KH7 ($10 \mu\text{M}$). KH7 was applied for ~ 30 minutes to maximise delivery and uptake into the CB before exposure to hypercapnia. **(e)** Mean single fibre discharge frequencies recorded in normocapnia ($\text{PCO}_2 = 40 \text{ mmHg}$) and hypercapnia ($\text{PCO}_2 = 80 \text{ mmHg}$), for control and during application of KH7. Error bars indicate \pm S.E.M. **(f)** Calculated CO_2 sensitivities ($\Delta \text{ Hz} / \text{ mmHg } \text{PCO}_2$) for control and KH7 groups. Error bars indicate \pm S.E.M. For E) and F) data is from 5 fibres ($n=5$) from 4 CB preparations.





